

Switchgrass Variety Choice In Europe¹

by H.W. Elbersen¹, D.G. Christian², N. El Bassam³, W. Bacher³, G. Sauerbeck³, E. Alexopoulou⁴, N. Sharma⁵, I. Piscioneri⁵, P. de Visser⁶ and D. van den Berg⁶

¹ATO (Agrotechnological Research Institute), Wageningen, NL e-mail: h.w.elbersen@ato.wag-ur.nl; ²IACR Rothamsted, United Kingdom, ³FAL (Federal Anstalt Fuer Landwirtschaft), Braunschweig, Germany; ⁴CRES (Center for Renewable Energy Sources), Pikermi, Attiki, Greece; ⁵ENEA (Ente per le Nuove Tecnologie, Energie e l'Ambiente), Italy; ⁶BTG (Biomass Technology Group), Enschede, The Netherlands.

Summary

Switchgrass is a perennial C₄ grass native to North America, where it occurs naturally from 55° N latitude to deep into Mexico. It is used for soil conservation, forage production, as an ornamental grass and more recently as a biomass crop for ethanol, fibre, electricity and heat production. In Europe research into the crop has just started and the choice of varieties for different geographical areas is an important issue. Some 20 different varieties have been evaluated for adaptation to different regions of Europe. The main factor determining area of adaptation of a variety is latitude of origin. Yields of varieties are correlated to the latitude of origin of the variety, with southern varieties having higher yield potential. If varieties are grown too far north they fail to winter-harden which decreases biomass quality (high nutrient and moisture content) and winter survival. It appears that in Europe switchgrass may be grown further north than in North America. The best variety for a given latitude or geographical area will be a compromise between yield and quality and long term winter survival.

Key words: Switchgrass, *Panicum virgatum*, quality, latitude, mineral composition.

Introduction

Switchgrass (*Panicum virgatum* L.) is a perennial C₄ grass native to North America where it occurs naturally from 55° N latitude to deep into Mexico, mostly as a prairie grass. In North America it has been used for more than 50 years for soil conservation, as a fodder crop and as an ornamental grass. Over the last two decades it has become an important warm-season pasture grass for fodder production when cool season C₃ grasses are less productive in summer (Moser and Vogel, 1995). Since the early 1990s the crop has been developed by the United States Department of Energy (DOE) as a model herbaceous energy crop for ethanol and electricity production. In Canada, Resource Efficient Agricultural Production (REAP) has

¹ REFERENCE: Elbersen, H. W., D. G. Christian, N. El Bassam, W. Bacher, G. Sauerbeck, E. Alexopoulou, N. Sharma, I. Piscioneri, P. De Visser, and D. Van Den Berg. 2001. Switchgrass variety choice in Europe. *Aspects of Applied Biology* 65: 21-28.

worked on switchgrass since 1991 for thermal conversion (electricity and heat) and ethanol production and is involved in projects to use switchgrass for paper pulp production. Many reasons are given for using switchgrass as a biomass crop for energy and fibre production, including the high net energy production per ha, low production costs, low nutrient requirement, low ash content, high water use efficiency, large range of geographic adaptation, ease of establishment by seed, adaptation to marginal soils, and potential for carbon storage in soil, (Christian and Elbersen, 1998; Samson and Omielan, 1992, Sanderson et al., 1996). In Europe research into the use as a biomass crop for energy and fibre has only just started. The crop has the potential to play a role in supporting policies to increase the use of durable products, reduce CO₂ emissions, utilise marginal and set aside lands and provide new economic activities for rural communities. Over the last years many larger and smaller individual field evaluations of switchgrass have been conducted (Christian and Elbersen, 1998; Lewandowski et al., 1998). We estimate that in Europe some 4 ha of experimental switchgrass fields exist of which 2.5 ha is within the current European Union sponsored switchgrass productivity network. In this network 6 organisations co-operate in evaluating the agronomic, fibre and energy potential of more than 20 switchgrass varieties under European conditions. The data from this project have been used in the current paper. Here we will discuss the important issue of allocation of varieties in relation to latitude of origin of the variety and the effect on establishment, yield and quality for energy production.

Available switchgrass material

Ecotypes

Two switchgrass ecotypes are generally defined based on morphological characteristics and habitat preferences. Lowland types are generally found in floodplains. They are taller, coarse, have a more bunch type growth habit, and may be more rapid growing than upland types. Upland types are found in drier upland sites. They are finer stemmed, and often semi-decumbent (Moser and Vogel, 1995; Porter, 1966). Artificial hybridization between lowlands and uplands have largely been unsuccessful (Taliaferro and Hopkins, 1997). Switchgrass is highly polymorphic and largely self incompatible (Talbert et al., 1983; Taliaferro and Hopkins, 1997). The basic chromosome number of switchgrass is $x = 9$. The ploidy levels of switchgrass range from diploid ($2n=18$) to duodecaploid ($2n=108$) (Hulquist et al., 1996; McMillan, 1959; Nielsen, 1944; Riley and Vogel, 1982). Most varieties are tetraploid or octoploid. Switchgrass varieties have been developed since the 1930s. In Table 1 the ploidy levels of available switchgrass varieties are given.

Varieties

Early varieties like Blackwell and Nebraska-28 are wild accessions that showed good performance and were released without additional breeding work (Moser and Vogel, 1995). The earlier varieties were mostly selected for soil conservation purposes, for example Dacotah and Alamo. More recent varieties have been developed by breeding for establishment, yield, quality, and disease resistance (Moser and Vogel, 1995; Sanderson et al., 1996). Currently breeding takes place at several locations in North America. New lowland and upland varieties are being developed specifically for biomass production for biofuel (Taliaferro and Hopkins,

1997). The high genetic variation for important traits should make development of improved varieties for several purposes possible.

Table 1. Ecotype, ploidy level, origin, and seed weight of available switchgrass varieties.

Variety	Ecotype	Ploidy level	Origin	Seed weight†
Alamo	lowland	Tetraploid	South Texas 28°	94
Blackwell	upland	Octoploid	Northern Oklahoma 37°	142
Caddo	upland	Octoploid	South Great plains 35°	159
Carthage = NJ-50	?	?	North Carolina 35°	148
Cave-in-Rock	Intermediate?	Octoploid	Southern Illinois 38°	166
Dacotah	upland	Tetraploid?	North Dakota 46°	148
Forestburg	upland	Tetraploid?	South Dakota 44°	146
Kanlow	lowland	Tetraploid	Central Oklahoma 35°	85
Nebraska 28	upland	?	Northern Nebraska 42°	162
Pangburn	lowland	Tetraploid	Arkansas 34°	96
Pathfinder	upland	Octoploid	Nebraska / Kansas 40°	187
REAP 921	upland	Tetraploid	Southern Nebraska 41°	90
Shelter = NY4006	mixed?	Octoploid?	West Virginia 40°	179
Summer	upland	Tetraploid	South Nebraska 41°	114
Sunburst	upland	?	South Dakota 44°	198
Trailblazer	upland	Octoploid	Nebraska 40°	185

(Alderson and Sharp, 1993; Anonymous, 1979; Barker et al., 1988; Barker et al., 1990; Boe and Ross, 1998; George and Reigh, 1987; Gunter et al., 1996; Hopkins et al., 1995; Hopkins et al., 1996; Jung et al., 1990; Newell, 1968; Stout et al., 1988; Vogel et al., 1991; Vogel et al., 1996). A question mark indicates that there are contradictions in the literature.

† Seed weight is expressed as seed weight per 100 seeds in mg. Reported seed weights are those found by the authors in one or two seed samples but should be typical for the variety.

Variety choice

Winter survival

It is known that varieties that are moved too far north from their origin will fail to mature in fall and have reduced winter survival. This is illustrated in Figure 1 for switchgrass varieties differing in latitude of origin grown in the Netherlands. Southern varieties did not flower in the fall and were still partially green when while northern varieties had flowered and were mature (brown). A frost killed the above ground (green) parts of the southern varieties in November. The failure to winter harden before a killing frost lead to reduced re-growth in spring (Figure 1.). The delayed re-growth can then have a negative effect on the yield as discussed below.

Yield

The most important factor determining area of adaptation of switchgrass varieties is latitude of origin. The plant has a photoperiod response which is modified by growing degree days (Moser and Vogel, 1995). Decreasing day-length will induce flowering in early summer. Other factors determining adaptation are precipitation and humidity. Varieties developed in dry areas will be more susceptible to fungal diseases when grown in humid conditions.




Figure 1. Relationship between latitude of origin of switchgrass varieties and maturity rating in fall and re-growth in early spring. First (establishment) year at Wageningen (NL).

When different varieties are grown at the same site northern ecotypes will remain shorter, flower earlier and mature earlier than southern ecotypes. Also, production of biomass will be considerably less compared to southern types (Jacobson et al., 1984). A clear strong correlation has been found (in N. America) between time to maturity, latitude of origin of the variety and yield (Sanderson et al., 1999, Samson et al., 1997). This effect can also be found in Europe as demonstrated in Figure 1A. Southern varieties matured later and had higher yields than northern varieties.

At northern sites (Figure 1B) intermediate varieties appeared to have highest yields in the second year. The lower yields for the southern varieties at Rothamsted (UK) and Noordoostpolder (NL) can be explained by lower winter survival and re-growth of southern varieties in the first (establishment) and subsequent years. Southern varieties mature too late to winter harden and translocate nutrients before winter sets in, leading to reduced re-growth in spring and lower yields. This is illustrated in Figure 2. (Figure 1B).

Quality

Quality of biomass for energy purposes depends on the type of application. Generally low moisture contents are required to reduce transportation costs and make storage possible. Furthermore low ash and nutrient contents are required when the biomass is used for combustion. K and Cl lower biomass quality by lowering the ash melting point and increasing corrosion problems in combustion systems. Nitrogen content increases NOx emissions.

Southern varieties that mature very late in fall will fail to translocate nutrients to the below ground parts before winter sets in. This is illustrated by the results of nutrient analysis in the Netherlands (Table 2) in the second year after establishment.




Figure 2. Relationship between latitude of origin of switchgrass varieties and second year biomass yields (A and B) and moisture content of the biomass (C,D). The experimental sites were located at 38° north (Aliartos, Greece), 40° north (Trisaia, Italy), 52° north (Rothamsted, UK) and 52° north (Wageningen, the Netherlands). Harvests were made in winter and early spring.

At a clay and a sandy site in two consecutive years varieties originating at northern latitudes (Forestburg) consistently had lower Cl and K contents than the southern variety Carthage while the intermediate varieties had intermediate Cl and K contents. N and P, nutrients which are not mobilised as easily, were also relatively low in the northern variety Forestburg but here the effects of latitude was less evident, especially in the 2000/2001 season. The same was the case for ash content. An explanation may lie in the heavy lodging of switchgrass, especially at the clay site, in 2000 which impeded trans-location of nutrients also for northern varieties. At the clay site much higher ash contents were measured as can be expected. This illustrates that apart from latitude of origin also other variables are important in determining biomass quality (and yield).

Thus it is likely that southern varieties require more fertiliser especially K and to a lesser extent N and P. Southern varieties generally have higher water contents at harvest because they mature later and have thicker stems. This reduces the harvest window and biomass quality (Christian and Elbersen, 1998). This is illustrated in Figure 1C and D, where northern varieties had lower moisture content at harvest both at southern and west European sites.

Table 2. Ash and nutrient content of switchgrass varieties differing in latitude of origin. The plots were established in 1998 on a clay soil type (Noordoostpolder) and on a sandy soil (Wageningen). The latitude of the two sites is approximately 52 and 53 °north. Plots were harvested after a killing frost in winter.

	-----Noordoostpolder, clay site -----						----- Wageningen, sandy site -----				
	Latitude	Ash	Cl	N	P	K	Ash	Cl	N	P	K
	°north	% DW	----- kg/tonne -----			% DW	----- kg/tonne -----				
1999-2000											
Forestburg	44	5,83	0,32	4,13	0,52	1,06	1,60	0,41	4,26	0,61	1,36
Summer	41	4,87	0,27	4,06	0,54	1,23	1,80	0,27	4,31	0,58	1,03
CIR	38	5,51	0,97	6,69	0,67	4,38	2,20	1,10	4,56	0,58	2,46
Blackwell	37	5,80	0,86	5,27	0,69	2,93	2,60	0,83	4,66	0,66	2,34
Carthage	35	5,30	1,48	6,56	0,88	5,61	2,60	0,83	5,86	0,75	3,31
2000-2001											
Forestburg	44	8,90	0,35	7,55	0,88	1,74	2,20	0,23	7,37	0,83	1,40
Summer	41	7,88	0,36	8,73	1,03	2,43	2,20	0,24	7,22	0,81	2,03
CIR	38	8,51	0,59	9,55	1,05	3,47	1,20	0,46	8,05	0,86	2,76
Blackwell	37	10,20	0,51	8,55	1,03	2,61	2,60	0,49	9,11	1,13	2,70
Carthage	35	8,50	0,54	8,83	1,12	3,35	2,00	0,52	7,39	0,99	2,72

Conclusions

It is possible to find switchgrass varieties that are adapted to most regions of Europe. The latitude of origin of a variety is the most important aspect determining the area of adaptation of a variety. Generally the use of varieties originating at southern latitudes can increase DM yields but it will also increase the chance of establishment failures in the first year, and a decline in yields over time. Furthermore the quality of the biomass will be reduced (high moisture and nutrient content) if the variety does not mature in the fall. The best variety for a given latitude or geographical area will be a compromise between yield, quality and winter survival. From the current data on switchgrass grown in Europe it appears that switchgrass may be grown further north than in North America.

Acknowledgements

This research has been partially funded by the European Community (DGVI) under the FAIR program (FAIR 5-CT97-3701). This paper does not necessarily reflect the European Commissions views or anticipate in any way its future policy in this area.

References

- Alderson, J., and W.C. Sharp. 1993.** Grass varieties in the United States, Agricultural Handbook No. 170. USDA, Washington D.C.
- Anonymous. 1979.** Alamo switchgrass. Bulletin L.1774. Texas Agricultural Experiment Station, College Station.
- Barker, R.E., R.J. Haas, E.T. Jacobson, and J.D. Berdahl. 1988.** Registration of 'Forestburg' switchgrass. Crop Sci. 28:192-193.

- Barker, R.E., R.J. Haas, J.D. Berdahl, and E.T. Jacobson. 1990.** Registration of 'Dacotah' switchgrass. *Crop Sci.*, 30:1158.
- Boe, A., and J.G. Ross. 1998.** Registration of 'Sunburst' switchgrass. *Crop Sci.*, 38:540.
- Christian D.G., and H.W. Elbersen. 1998. Switchgrass (*Panicum virgatum* L.). In: N. El Bassam. Energy plant species. Their use and impact on environment and development. London: James and James publishers, p. 257-263.
- George, J.R., and G.S. Reigh. 1987.** Spring growth and tiller characteristics of switchgrass. *Canadian Journal of Plant Science*, 67:167-174.
- Gunter, L.E., G.A. Tuskan, and S.D. Wullschleger. 1996.** Diversity among populations of switchgrass based on RAPD markers. *Crop Sci.*, 36:1017-1022.
- Hopkins, A.A., K.P. Vogel, K.J. Moore, K.D. Johnson, and I.T. Carlson. 1995.** Genotype effects and genotype by environment interactions for traits of elite switchgrass populations. *Crop Sci.*, 35:125-132.
- Hopkins, A.A., C.M. Taliaferro, C.D. Murphy, and D.A. Christian. 1996.** Chromosome number and nuclear DNA content of several switchgrass populations. *Crop Sci.*, 36:1192-1195.
- Hulquist, S.J., K.P. Vogel, D.J. Lee, K. Arumuganathan, and S. Kaeppler. 1996.** Chloroplast DNA and nuclear DNA content variations among cultivars of switchgrass, *Panicum virgatum* L. *Crop Sci.* 36: 1049-1052.
- Jacobson, E.T., D.A. Tober, R.J. Haas, and D.C. Darris. 1984.** The performance of selected cultivars of warm season grasses in the northern prairie and plains States. In: G. K. Clambey and R. H. Pemble (eds.) *The Prairie: Past, present and future*, 9 ed, Moorhead Minnesota. pp. 215-226.
- Jung, G.A., J.A. Shaffer, W.L. Stout, and M.T. Panciera. 1990.** Warm-season grass diversity in yield, plant morphology, and nitrogen concentration and removal in Northeastern USA. *Agron. J.*, 82, 21-26.
- Lewandowski, I., R. Olsson, W. Bacher, W. Huisman, L.S. Nikolaisen, and L. Paavilainen. 1998.** Production and use of perennial rhizomatous grasses (PRG) in the energy and industrial sector of Europe. In: I. Lewandowski (ed.) *Workshop at the 10th European conference on "Biomass for Energy and Industry"*, vol. 1, Wurzburg, Germany. pp. 80.
- McMillan, C. 1959.** The role of ecotypic variation in the distribution of the central grasslands of North America. *Ecol. Monogr.* 29: 285-308.
- Moser, L.E., and K.P. Vogel. 1995.** Switchgrass, Big Bluestem, and Indiangrass. In: *An introduction to grassland agriculture*. R. F. Barnes, D. A. Miller and C. J. Nelson (eds.). Forages, 5th ed. Vol.1, Ames, IA: Iowa University Press, pp. 409-420.
- Newell, L. C. 1968.** Registration of Pathfinder switchgrass. *Crop Science* 8, 516.
- Nielsen, E.L. 1944.** Analysis of variation in *Panicum virgatum*. *J. Agric. Res.* 69: 327-353.
- Porter C.L. 1966. An analysis of variation between upland and lowland switchgrass, *Panicum virgatum* L., in central Oklahoma. *Ecology*, 47:980-992.
- Riley, R.D. and K.P. Vogel. 1982.** Chromosome numbers of released cultivars of switchgrass, indiangrass, big bluestem, and sand bluestem. *Crop Sci.* 22: 1082-1083.
- Samson, R.A., and J.A. Omielan. 1992.** Switchgrass: A potential biomass energy crop for ethanol production. In: *The Thirteenth North American Prairie conference*, Windsor, Ontario. p. 253-258.
- Samson, R., P. Girouard, and Y. Chen. 1997.** Evaluation of switchgrass and short rotation forestry willow in eastern Canada as bio-energy and agri-fibre feedstocks. In: R. P. Overend and E. Chornet (eds.) *Proceedings of the third conference of the Americas*, Montreal, Canada. pp. 145-151.

- Sanderson, M.A., R.L. Reed, S.B. McLaughlin, S.D. Wullschleger, B.V. Conger, D.J. Parrish, D.D. Wolf, C. Taliaferro, A.A. Hopkins, W.R. Ocumpaugh, M.A. Hussey, J.C. Read, and C.R. Tischler. 1996.** Switchgrass as a sustainable bioenergy crop. *Bioresource Technology*. 56: 83-93.
- Sanderson, M.A., R.L. Reed, W.R. Ocumpaugh, M.A. Hussey, G.v. Esbroeck, J.C. Read, C.R. Tischler, F.M. Hons. 1999.** Switchgrass cultivars and germplasm for biomass feedstock production in Texas. *Bioresource Technology*. 67: 209-219.
- Stout, W.L., G.A. Jung, and J.A. Shaffer. 1988.** Effects of soil and nitrogen on water use efficiency of tall fescue and switchgrass under humid conditions. *Soil Science Society of America Journal*, 52:429-434.
- Talbert, L.E., D.H. Timothy, J.C. Burns, J.O. Rawlings and R.H. Moll. 1983.** Estimates of genetic parameters in switchgrass. *Crop Sci*. 23: 725-728.
- Taliaferro, C.M. and A.A. Hopkins. 1997.** Breeding and selecting of new switchgrass varieties for increased biomass production. Five year summary report. Oak Ridge National Laboratory, Oak Ridge, TN. contact #19XSL127C.
- Vogel, K.P., F.A. Haskins, H.J. Gorz, B.A. Anderson, and J.K. Ward. 1991.** Registration of 'Trailblazer' switchgrass. *Crop Sci.*, 31:1388.
- Vogel, K.P., A.A Hopkins, K.J. Moore, K.D. Johnson, and I.T. Carlson. 1996.** Registration of 'Shawnee' switchgrass. *Crop Sci.*, 36:1713.